

VISIBLE LIGHT



These views of the constellation Orion dramatically illustrate the difference between the familiar, visible-light view and the richness of the universe that is invisible to our eyes, though accessible via other parts of the electromagnetic spectrum.



Infrared Astronomy: More Than Our Eyes Can See

Infrared Astronomy: More Than Our Eyes Can See

Invisible Light. In 1800, William Herschel discovered an invisible form of radiation just beyond the red portion of the visible spectrum. He named this form of radiation infrared ("below" red). (Herschel was already world-famous for having discovered the planet Uranus.) Herschel's discovery was the first step in establishing the existence of what we now call the electromagnetic spectrum. Visible light and infrared radiation are just two of the many types of electromagnetic energy produced by objects on Earth and throughout the universe. Only by studying all of these types of radiation can we fully characterize celestial objects and gain a complete picture of the universe, its history, and its evolution.

Studying the Hidden Universe. The heat that we feel from our Sun or from a fireplace is infrared (sometimes called thermal) radiation. Even objects that we think of as being very cold, such as ice cubes, emit infrared radiation. Measuring the infrared energy arriving from astronomical objects is difficult because much of it is blocked by water vapor and other molecules in Earth's atmosphere. Therefore, most infrared astronomers use airborne telescopes, balloon payloads or space telescopes to study the thermal radiation from celestial objects. The telescopes and detectors used by infrared astronomers emit their own infrared radiation. To minimize this interference, and to be able to detect the much weaker radiation from celestial objects of interest, infrared astronomers refrigerate their instruments and even entire telescopes to temperatures as cold as -269 °C (-452 °F) — almost absolute zero!

Building New Stars. The visible and infrared images shown on the other side of this lithograph are of exactly the same region of the sky containing the constellation Orion. These images dramatically illustrate how features that cannot be seen in visible light show up very brightly in the infrared. The infrared image shows several regions of hot, dense cores within clouds of gas and dust. These are the stellar nurseries where new stars like the Sun and planets like the Earth are being born. Embedded in these molecular clouds, the young stars are difficult to see in visible light, but their presence is revealed by infrared radiation.

The interstellar medium (ISM) in our Milky Way galaxy is the dust and gas between the stars. Some of it is primordial, dating back to the origin of the universe, and some has been added by the violent deaths of massive stars in supernova explosions, or in gentler episodes when the outer layers of some stars are blown off near the ends of their lives. The ISM is the reservoir of material from which new stars form. Clouds of interstellar dust and gas are easily detected at long infrared wavelengths (about 100 times longer than visible radiation). To see the newborn stars embedded in these clouds of dust and gas, astronomers depend instead on observations made at short infrared wavelengths that penetrate the obscuring dust.

Building New Planets. In the 1980s and 1990s, using data from the *Infrared Astronomical Satellite (IRAS)* and the *Hubble Space Telescope (HST)*, astronomers discovered disks of dust around over a hundred nearby stars. These disks contain the raw materials from which planetary solar systems are formed: as such, these disks provided the first tantalizing evidence that planets orbiting stars are probably common occurrences.

Understanding Galaxies. Our view of distant regions of the Milky Way galaxy, including the galactic center, is heavily obscured by the ISM at wavelengths of visible light. The galactic center is one of the brightest sources of infrared radiation in the sky. Infrared observations show that this region consists of very dense crowds of stars and clouds of gas which orbit the galactic center unexpectedly rapidly — evidence of the gravitational influence of a massive central black hole.

Observing the Past. When we observe galaxies at distances of billions of light years, we see them as they existed billions of years ago because of the finite speed of light. Moreover, the universe is known to be expanding, producing a shift in the measured wavelength of radiation relative to the wavelength at which it was emitted. This shift of the visible radiation toward the red end of the electromagnetic spectrum is known as the "cosmological red shift." If the emitting objects are sufficiently far away (long ago), the radiation received is shifted from the visible spectrum into the infrared. Thus, infrared observations provide a glimpse of the early universe, an era when the first stars and galaxies were forming.

NASA Infrared Astronomy Missions

• The Spitzer Space Telescope completed NASA's set of multi-wavelength "Great Observatories" which also included the Hubble Space Telescope, the Compton Gamma Ray Observatory, and the Chandra X-ray Observatory. Spitzer studied the Universe at near-, mid-, and far-IR wavelengths with an 85 cm (33 in.) diameter telescope in solar orbit during the 5-1/2 year lifetime of its cryogen supply. Spitzer continued to operate part of its near-IR camera for an additional 11 vears. • The Stratospheric Observatory for Infrared Astronomy (SOFIA) was a Boeing 747SP airplane modified to carry a telescope with a diameter of 2.7 m (106 in.), a joint project of NASA and the German Aerospace Center, DLR. Flying at altitudes of 12-14 km (39,000-45,000 ft), SOFIA operated above most of our atmosphere's obscuring water vapor, concentrating on the far-IR and sub-millimeter wavelength ranges. SOFIA's scientific instruments were especially well suited for comparing how stars form in our Milky Way Galaxy versus other galaxies, understanding how magnetic fields affect the interstellar medium, and investigating the production of organic compounds in space. The SOFIA mission ended in 2022.

• The Wide-field Infrared Survey Explorer (WISE) was a NASA space telescope mission launched in 2009 that operated in Earth orbit for almost a year until its cryogen supply was depleted. Its 40 cm (16 in.) telescope mapped the entire sky at near- and mid-IR wavelengths with 500 times greater sensitivity than previous space-borne infrared survey missions. Astronomers continue to analyze WISE data to study the nearest and coolest stars, the origins of stellar and planetary systems, and the most luminous galaxies in the Universe. After a 2-year hiatus, WISE was re-activated for 11 years as NEOWISE, using its near-IR detectors to search for Near Earth Objects (NEOs) until the mission ended in 2024.

• The James Webb Space Telescope (JWST) is a space observatory launched into solar orbit in 2021 by NASA in partnership with the European and Canadian space agencies. JWST has a 6.5 m (256 in.) diameter primary mirror providing unprecedented sensitivity and spatial resolution in the near- and mid-IR wavelength ranges. Two of JWST's key projects are to study the formation of galaxies in the early Universe and the composition of extrasolar planet atmospheres and surfaces.

For more information:

https://science.nasa.gov/mission/spitzer/ (Spitzer Space Telescope) https://science.nasa.gov/mission/sofia/ (SOFIA) https://science.nasa.gov/mission/neowise/ (WISE/NEOWISE) https://science.nasa.gov/mission/webb/ (James Webb Space Telescope) https://coolcosmos.ipac.caltech.edu (infrared light and infrared astronomy tutorial)

Photo Credits: Visible-light image, Akito Fujii/David Malin Images; Infrared image, NASA/IRAS/Deborah Levine

https://science.nasa.gov/learn/

NASA SMD SciAct

Astronomy Activation Ambassadors

https://science.nasa.gov/sciact-team/astronomy-activation-ambassadors/ https://www.seti.org/aaa